

ATROPINE INJECTION, BP – AMPOULE

Class	Anticholinergic and Antimuscarinic
Description	Solution does not contain any antimicrobial agent, or bacteriostat. It is intended as a single dose ampoule. When smaller dosage is needed, remaining solution should be discarded
Contra-indications	Atropine should not be given to patients with closed-angle glaucoma, with myasthenia gravis, with prostatic enlargement. Atropine should not be given to patients who are being treated with a monoamine-oxidase inhibitor, or within ten days of the discontinuation of such treatment
Indications	Before the induction of general anaesthesia. In patients with an acute myocardial infarction who have developed a bradycardia or atrioventricular block. For the treatment of intoxication with cholinesterase inhibitors and for intoxication by Amanita muscaria and other mushroom poisonings where the toxic agent is a muscarine-like compound
Dosage	Adults: Premedication, IV 0.3-0.6mg immediately before induction of anaesthesia and incremental doses of 0.1mg for the treatment of bradycardia – IM 0.3-0.6mg 30-60 minutes before induction.

SUPPLY SPECIFICAITONS

Product No	Description	Strength	Size	Unit of Sale
431	Single Dose Ampoule	0.6mg/mL	1ml	100

COMPOSITION

Atropine	0.6mg/mL
Sodium Chloride	To render solution isotonic
In Water for injection	0ml

**Information provided for marketing purpose only.
Consult physician for proper administration.**